

CLUMPUSCULE FORMATION AT HIGH REDSHIFT

F. COMBES

*Observatoire de Paris, DEMIRM
61 Av. de l'Observatoire, F-75 014 Paris, France*

and

D. PFENNIGER

*Observatoire et Université de Genève
CH-1290 Sauverny, Switzerland*

ABSTRACT. Over the last two decades, realistic studies have often conclude that the first bound objects to form can be very small, much smaller than a solar mass. After recombination, the Jeans mass drops rapidly to the order of a Giant Molecular Cloud (GMC) mass ($\sim 10^5 M_\odot$), and the H_2 cooling can make the collapse quasi-isothermal; this leads to recursive fragmentation, and formation of clumps so dense that 3-body reactions transform the gas almost entirely to the molecular phase. This could lead to star formation in some places, but since star formation is very inefficient, most of the molecular gas could consist of a fractal built on clumpuscules thermalized with the background radiation, and filling a tiny fraction of the volume. *The bulk of the gas mass can therefore be trapped in this phase, well before the first stars re-heat and re-ionize the diffuse gaseous medium.* This results in a very contrasted multi-phase baryonic medium, that has partly remained until the present time.

1. The first bound structures

1.1. General background

In standard big-bang cosmologies, observable structures are assumed to form from small density fluctuations, that first expand with the Universe while growing linearly, then collapse non-linearly to bound systems. These fluctuations could be adiabatic, in which case they would be damped on scales below $3 \cdot 10^{13} (\Omega h^2)^{-5/4} M_\odot$ before recombination (Silk 1968; where h is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$). Isothermal fluctuations can also exist (and, of course, also intermediate states), and in such cases, the first non-linear objects are much smaller. If the density fluctuations spectrum is assumed self-similar, with a slope n such that

$$\frac{\delta\rho}{\rho} \propto M^{-(n+3)/6} \quad (1)$$

then the largest masses to become non-linear after recombination, i.e. at $z \sim 1500$, are of the order of $M \sim 10^{6-8} M_\odot$, depending on Ω , h and n , if the spectrum is calibrated from the presently observed correlation function (e.g., Peebles 1980). That means that all masses smaller than $M \sim 10^{6-8} M_\odot$, and larger than the Jeans mass, which is $M \sim 10^5 M_\odot (\frac{\Omega_b}{0.06})^{-1/2} (\frac{h}{0.5})^{-1}$ at $z = 1000$, will collapse, become bound, and decouple

from expansion (at this epoch the Jeans length is of the order of $2 \text{ pc } (\frac{\Omega_b}{0.06})^{-1/2} (\frac{h}{0.5})^{-1}$). In fact, since in every hierarchical model with $n > -3$ the smallest masses have more density contrast, structures at precisely the Jeans mass collapse first. The masses correspond to typical $z = 0$ giant molecular clouds.

The nature of the collapse has been studied by many authors. If cooling is efficient enough ($\tau_{\text{cool}} \sim \tau_{\text{ff}}$), the collapse is quasi-isothermal, and fragmentation occurs, since the Jeans mass becomes smaller and smaller as the density increases (e.g., Hoyle 1953). This means that pressure forces are at most equal to gravity forces, and cannot stop the collapse. This fragmentation process occurs currently in the Galactic interstellar medium in a self-similar hierarchical structure (Larson 1981; Scalo 1985), and might even be pursued down to very low masses in particularly cold and quiet regions, such as the outer parts of galaxies (Pfenniger & Combes 1994). Fragmentation is limited by opacity, and the smallest fragments (or clumpuscles), which are at the transition of being pressure supported, are today of the order of $10^{-3} M_{\odot}$, and their mass grows slowly, as $T^{1/4}$ or $(1+z)^{1/4}$, with redshift. This assumes a quasi-isothermal regime for the bottom of the hierarchy (containing most of the mass); therefore it is important that gas cooling remains efficient.

This might look as a problem for gas just after recombination, since it is metal poor without grains: if above 10^4 K the main coolant is atomic hydrogen (by collisional excitation of $\text{Ly}\alpha$), in fact below this temperature the vibration-rotation lines of molecular hydrogen take over until $T = 200 \text{ K}$, because a significant quantity of H_2 molecules forms through H^- and H_2^+ . Below 200 K , HD is then more efficient (Palla & Zinnecker 1987; Puy & Signore 1996). Many groups have tackled the problem of computing the physico-chemistry of the primordial gas, to determine the size of the first forming bound structures (Yoneyama 1972; Hutchins 1976; Carlberg 1981; Palla et al. 1983; Lepp & Shull 1984). All of them have found that the cooling is indeed efficient, as soon as the redshift is below $z \sim 200$, and the calculated masses of fragments are in the wide range $0.1 - 100 M_{\odot}$. Even if clouds of just the Jeans mass see their collapse somewhat delayed because of pressure forces, the latter are negligible for clouds of $M > 10^6 M_{\odot}$ (Lahav 1986). In more realistic scenarios, the collapse is not spherical, but does form sheets and filaments (Larson 1985). The latter have the advantage that in 1D, pressure forces have always the same dependence with radius than the gravity forces, and therefore cannot halt the collapse, whatever the cooling (Uehara et al. 1996; Inutsuka & Miyama 1997). Therefore fragmentation is inevitable.

The conclusions drawn from these studies were all concerning the first generation of stars, since *no outcome to fragmentation other than star formation was considered*. Already Peebles & Dicke (1968) concluded that the first objects to form were the globular clusters, through such a fragmentation process, followed by star formation. Palla et al. (1983) considered the H_2 formation through the very efficient 3-body process, when the density is higher than 10^8 cm^{-3} . They concluded that all the primordial gas is converted in H_2 at a density of 10^{12} cm^{-3} , and that the Jeans mass eventually falls below $0.1 M_{\odot}$, allowing the whole mass spectrum for the first forming stars. Rotation could hinder the collapse (Kashlinsky & Rees 1983), and this could lead to Population III low-mass stars as well as super-massive objects (VMO's), both of which could contribute to dark matter.

Taking the observations of the ISM at face, which show that the efficiency of star formation is very low, we propose here that the bulk of the gas in this first collapse of GMC-size primordial clouds after recombination and decoupling form clumpuscles, assembled in a hierarchical or fractal structure. The molecular density of these clumpuscles should be very high (10^{10} to 10^{12} cm^{-3}), when at the limit of being pressure supported, at the temperature of the cosmic background.

1.2. Non star-formation

Although theoretically not well understood, observations show that the efficiency of star formation is highly variable, and must be very low in some cases (e.g., in low surface brightness galaxies such as Malin 1). Since the physical conditions of the gas at $z \sim 50 - 100$ is similar to the present outer parts of galaxies, we expect that the conditions for triggering global star formation are generally not met. Larger scales structures such as the galaxy-size ones, have not then collapsed and decoupled from expansion, therefore there does not exist a deep enough potential well to accumulate mass to the critical density for star-formation (Kennicutt 1989). Indeed, star formation is hindered when the medium is only mildly unstable, so that pressure forces compensate gravity forces. That is the case, when the collapsing masses are always of the order of the Jeans mass. In the first collapse of the giant molecular clouds of $10^6 M_{\odot}$, this is the case, since the Jeans mass decreases gradually as fragmentation proceeds. When galaxy sized masses collapse, leading to more violent instabilities, large fluctuations, and shocks at the origin of starbursts, they are much higher than the Jeans mass. However, when a disk forms, rotation takes over, and also compensates for the gravity forces at large scale. The pressure forces stabilise the smallest scales (given by the Toomre criterion), and rotation stabilises the large scales (allowing disk structures to survive). Apparently, in galactic disks star formation occurs substantially when the stability conditions are violated at intermediate scales. Note that the Toomre criterion does not prevent the formation of substructures when the medium is rapidly cooling. We expect that the smallest fragments are in quasi-thermal equilibrium with the microwave background, and statistical dynamical equilibrium between coalescence and fragmentation over the whole hierarchy of structures (Pfenniger & Combes 1994). The dissipation is thus minimised, since the temperature of the fragments is very close to that of the cosmic background. Clumpuscles are thought as dynamical entities, they are frequently reshuffled by collisional disruption and coalescence. They permanently reform through Jeans instability and fragmentation, so the dynamical quasi-equilibrium is statistical along the whole hierarchy of clumps.

At high redshift ($z > 50 - 100$) therefore, only sporadic star-formation (or even MACHOS formation) should occur in rare places, and that might be sufficient to initiate the reionization and the re-heating of the intergalactic medium (IGM), as is necessary to explain the high ionization fraction of the Lyman- α absorbers (Tegmark et al. 1984) and the omnipresent HeII gas at $z \sim 2$ (Jacobsen et al. 1994; Davidsen et al. 1996; Reimers et al. 1997). The first generation of stars could form today a small percentage of the dark halos around galaxies under the form of white or brown dwarfs (Alcock et al. 1997).

1.3. Clumpuscles resistance to reionization

Once the bulk of the gas mass has condensed into cold molecular hydrogen at $z \sim 150$, the clumps survive the reionization phase of the IGM in large part, because of their high column density ($> 10^{25} \text{ cm}^{-2}$), that self-shields them from the radiation, and their fractal structure, which is more gregarious than an homogeneous distribution in space (cf. Combes & Pflamminger 1997). Some erosion occurs, mainly at the interfaces, which produces first atomic hydrogen, then ionized gas. But in the densest environments, that will become the galaxies, the bulk of the gas remains in cold clumpuscles. The estimation of the column density threshold of atomic hydrogen at the ionizing limit can be obtained from the measured and modeled values of the extra-galactic ionizing radiation, that is known now within a factor of a few. The extra-galactic background, essentially from the quasar UV light, provides an ionization rate of $\xi \sim 2 \cdot 10^{-14} \text{ s}^{-1}$, a value corresponding to the study of low-redshift Lyman- α absorption lines (Madau 1992). As in the “Strömgren sphere theory”, there is a sharp transition (a true ionizing front) between the neutral and ionized phases, and the transition limit is around $N_{\text{HI}} \sim 10^{18} \text{ cm}^{-2}$, for usual disk gas volumic densities n (it decreases as $1/n$). This corresponds well to the observed sudden decrease from a column density of $N_{\text{HI}} = 2 \cdot 10^{19} \text{ cm}^{-2}$ to $2 \cdot 10^{18} \text{ cm}^{-2}$ in the outskirts of HI disks (Corbelli & Salpeter 1993). Although the extra-galactic flux might increase with redshift, it seems therefore quite easy to conserve the cold H_2 clumps within their HI interface at high redshifts.

2. H_2 cooling

The details of H_2 chemistry and cooling has been widely developed (Lepp & Shull 1984; Haiman et al. 1996a; Tegmark et al. 1997). Numerical 3D simulations have been performed (Abel et al. 1997; Anninos et al. 1997), in the aim of determining the mass of the first luminous objects to form, and of following the effective Jeans mass as a function of redshift. It was shown that very small masses ($< 0.1 M_{\odot}$) could be formed before reionization ($z > 50 - 100$), but then the first stellar formation would produce enough UV radiation to destroy the molecules (Haiman et al. 1997), although the exact processes are complex (the UV irradiation could even trigger extra-cooling by favoring H_2 formation, Haiman et al. 1996b). This H_2 destruction through the reionization period is used to delay the formation of dwarf galaxies, since giant molecular clouds will not form again until low redshifts, when efficient cooling is provided by the metals (Norman & Spaans 1997; Kepner et al. 1997).

The formation of small molecular clouds before reionization is therefore not in doubt, and we show in a simple model resolving the chemistry and cooling equations, that high enough densities are indeed achieved. For the sake of comparison, we adopt a simple scheme very similar to that of Tegmark et al. (1997): since we stop when the clumpuscles become optically thick with respect to the H_2 rotational lines (i.e., at column densities of the order of $10^{24-25} \text{ cm}^{-2}$, and volumic density of $\sim 10^{10} \text{ cm}^{-3}$), we have to consider the gas thermal history of a unit volume only, irrespective of the mass of the condensed object. The density before virialisation is determined by the simple top-hat model, and when virialised, we consider its possible recursive fragmentation.




Fig. 1. Evolution of temperature (T_6 in 10^6 K), molecular fraction (f_{H_2}), ionization fraction (x_e) and density (n_3 in 10^3 cm^{-3}) for a gas cloud after recombination, collapsing at $z_{\text{vir}} = 100$ (left), and $z_{\text{vir}} = 50$ (right). The Hubble constant is $h = 0.5$, and $\Omega = \Omega_b = 0.06$.

To fix ideas, we adopt physical parameters compatible with the observed fractal of the Milky Way molecular ISM, a fractal dimension of $D = 1.7$, and a number of fragments of $N = 8$ at each level (Pfenniger & Combes 1994). The ratio between the different scales of the hierarchy is then $N^{1/D} = 3.4$ and the density ratio is $r_d = N^{(3-D)/D} = 4.9$. Assuming that the cooling is always efficient enough to evacuate the gravitational energy of the collapse, an hypothesis that will be confirmed a posteriori according to the virial masses considered, we can compute the density as a function of time by steps corresponding to the free-fall time $\tau_{\text{ff}i}$ of a given level i of the hierarchy: $n(t) = n_i r_d^{(t-t_i)/\tau_{\text{ff}i}}$. The H_2 molecules are formed essentially through the $\text{H}^+ + \text{H}$ (via H_2^+) reaction at high redshift ($z \sim 1000$, Shapiro & Kang 1987), then through the $\text{H}^- + \text{H}$ reaction at $z \sim 100$ (Hutchins 1976), and through the 3-body reactions ($\text{H} + \text{H} + \text{H}$) at very high densities (Palla et al. 1983). The cooling at $T < 1000 \text{ K}$ is essentially due to H_2 . At low temperature ($T < 100 \text{ K}$), the HD molecules can also intervene. The LiH molecules are formed in too low quantities to be helpful (Stancil et al. 1996). In any case, this occurs at so high column densities that the H_2 rotational lines are optically thick, which is not considered here.

Some results of the computations are displayed in Fig. 1, for $z_{\text{vir}} = 100$ and 50. The horizontal full line indicates the virial temperature of the Jeans mass at recombination, i.e., $M = 10^5 M_\odot$. At $z_{\text{vir}} = 100$, the computed gas temperature is always lower than T_{vir} , and therefore the hypothesis of fragmentation is verified for all masses able to collapse. Computations as a function of z_{vir} indicate that this is true for $z_{\text{vir}} > 60$. For lower z_{vir} , only larger masses would fragment down the whole hierarchy, unless

some metal enrichment has occurred through sporadic star formation before. *Therefore we have shown that a different hypothesis about the outcome of fragmentation leads to completely different results from Tegmark et al. (1997) concerning the amount of molecular hydrogen and the state of the bulk of the gas at redshifts < 50 .*

3. The self-gravitating fractal gas

3.1. Merging tendency

Before significant star formation has occurred in the cold gas component, the medium is still dissipative and self-gravitating, and ideas on quasi-isothermal fragmentation can be applied (de Vega et al. 1996). The stability against sub-clumps merging ensues. This is opposite to the behaviour of dissipationless components, such as MACHOS, stars or non-baryonic dark matter. In the latter case, it is assumed that as soon as a structure decouples from expansion, all the sub-structures already decoupled are washed out in a dynamical time through merging. This is due to violent relaxation and dynamical friction, that transfer the energy of relative orbital motions to the sub-structures themselves, and disrupt them (e.g., White & Rees 1978). On the contrary, in a dissipative collapse and fragmentation process, as soon as sub-structures coalesce, they form an entity larger than the Jeans mass, and can re-fragment in a free-fall time, or dynamical time for this scale (e.g., Pfenniger & Combes 1994). The fragmentation along the whole hierarchy of the Jeans mass after recombination is almost instantaneous, as soon as the Jeans mass has collapsed, as already noted by Hoyle (1953). The whole process takes only a small fraction of the collapse time-scale of the GMC at this epoch of $\tau \sim 1$ Myr.

3.2. Galaxy formation

After recombination and re-heating, the formation of larger and larger objects is possible as long as the corresponding fluctuations become non-linear. The formation of galaxies could range from redshifts $z = 50 - 100$ to the present according to their mass (at redshift $z = 200$ normal 10 kpc-radius galaxies would overlap, cf. Peebles 1980). Since the proto-galaxies contained mainly gas at the beginning, dissipation forms disks, and rotation support is provided by the angular momentum gained in tidal interactions. The proto-galaxy collapse is therefore a rather violent relaxation process, and the rate of star formation could be high (and higher in the largest mass objects). In any case, the relaxation is much shorter and violent in the center of the collapsed objects. In the outer parts, most of the mass can remain there in the form of self-gravitating cold gas, that has subsisted from its formation period before the reionization.

When a substantial part of the mass has formed a stellar dissipationless component, the galaxy population enters the phase of hierarchical merging. According to their environment, and their tidal interactions with companions, galaxies evolve at different speeds toward the early-type side of the Hubble sequence (Pfenniger, Combes & Martinet 1994). When structures of the group-size turn around, further stellar activity peaks occur. Later, the formation of clusters trigger again enhanced activity.

3.3. Cluster formation

Clusters form around redshift $z \sim 2$, according to their mass, and to the cosmological model adopted. Many are still assembling today, as is attested by the substructures observed. What happens then to the extended gas halos that still surround late-type galaxies? They are very likely to be stripped, either through tidal interactions, or to ram-pressure when a substantial and dense hot intergalactic gas component has formed. Already in the HI tracer, galaxies appear to be stripped in the center of clusters like Virgo and Coma (e.g. Cayatte et al. 1990). Some of the stripped gas is heated at the virial temperature of the cluster, and accounts for the huge amounts of gas mass detected in X-rays (e.g. David et al. 1995). But all the clumpuscules are not destroyed, and a multi-phase medium could survive in the IGM (Ferland et al. 1994), especially at the center of the cluster, where the cooling time is shorter than the dynamical time (cooling flows), and the cold clumpuscules can reform. *Therefore the bulk of the baryonic mass detected in clusters in the form of X-ray gas is probably only a lower limit to the existing gas, and therefore to the amount of baryons.*

4. Conclusions

We propose that just after recombination and decoupling, gaseous structures having the Jeans mass, of the order of a giant molecular cloud of $10^{5-6} M_{\odot}$, collapse and fragment in the same time to form cold molecular clumpuscules. The fragmentation can proceed to masses down to Jupiter masses, since the collapse is quasi-isothermal, due to the H_2 cooling. The large majority of these pressure-supported fragments do not form stars, since they are in statistical equilibrium between coalescence and fragmentation, in a fractal structure, and in thermal equilibrium with the cosmic background temperature. A very low level, sporadic, star formation is sufficient to re-ionize the intergalactic gas at $z = 50 - 100$. The bulk of the clumpuscules survive the reionization, and may be assembled later on in larger structures, when proto-galaxies form. This scenario provides a way to avoid the “cooling” catastrophe at large redshifts, that will produce much more stars that are observed today (Blanchard et al. 1992). It implies the existence of a large fraction of the dark matter around galaxies under the form of cold molecular gas (also consistent with the recent X-ray results in clusters, David 1997).

References

- Abel T., Anninos P., Zhang Y., Norman M.L.: 1997, *New Astron.* **2**, 181
- Alcock C. et al: 1997, *Astrophys. J.* **486**, 697
- Anninos P., Zhang Y., Abel T., Norman M.L.: 1997, *New Astron.* **2**, 209
- Blanchard A., Valls-Gabaud D., Mamon G.A.: 1992, *Astron. Astrophys.* **264**, 365
- Carlberg R.G.: 1981, *Mon. Not. R. Astr. Soc.* **197**, 1021
- Cayatte V., Balkowski C., van Gorkom J.H., Kotanyi C.: 1990, *Astron. J.* **100**, 604
- Combes F., Pfenniger D.: 1997, *Astron. Astrophys.* **327**, 453
- Corbelli E., Salpeter E.E.: 1993, *Astrophys. J.* **419**, 94-104
- David L.P.: 1997, *Astrophys. J.* **484**, L11
- David L.P., Jones C., Forman W.: 1995, *Astrophys. J.* **445**, 578

- Davidsen A.F., Kriss G.A., Zheng W.: 1996, *Nature* **380**, 47
 De Vega H., Sanchez N., Combes F.: 1996, *Nature* **383**, 56
 Ferland G.J., Fabian A.C., Johnstone R.M.: 1994, *Mon. Not. R. Astr. Soc.* **266**, 399
 Haiman Z., Thoul A.A., Loeb A.: 1996a, *Astrophys. J.* **464**, 523
 Haiman Z., Rees M.J., Loeb A.: 1996b, *Astrophys. J.* **467**, 522
 Haiman Z., Rees M.J., Loeb A.: 1997, *Astrophys. J.* **476**, 458
 Hoyle F.: 1953, *Astrophys. J.* **118**, 513
 Hutchins J.B.: 1976, *Astrophys. J.* **205**, 103
 Inutsuka S-I., Miyama S.M.: 1997, *Astrophys. J.* **480**, 681
 Jacobsen P. et al: 1994, *Nature* **370**, 35
 Kashlinsky A., Rees M.J.: 1983, *Mon. Not. R. Astr. Soc.* **205**, 955
 Kepner J.V., Babul A., Spergel D.N.: 1997, *Astrophys. J.* **487**, 61
 Kennicutt R.C.: 1989, *Astrophys. J.* **344**, 685
 Lahav O.: 1986, *Mon. Not. R. Astr. Soc.* **220**, 259
 Larson R.B.: 1981, *Mon. Not. R. Astr. Soc.* **194**, 809
 Larson R.B.: 1985, *Mon. Not. R. Astr. Soc.* **214**, 379
 Lepp S., Schull J.M.: 1984, *Astrophys. J.* **280**, 465
 Madau P. 1992, *Astrophys. J.* **389**, L1
 Norman C.A., Spaans M.: 1997, *Astrophys. J.* **480**, 145
 Palla F., Salpeter E.E., Stahler S.W.: 1983, *Astrophys. J.* **271**, 632
 Palla F., Zinnecker H.: 1987, in *Starbursts and Galaxy Evolution XXII* Ind Rencontres de Moriond, ed. T.X. Thuan, T. Montmerle & J.T.T. Van Ed. Frontières, page 533
 Peebles P.J.E., Dicke R.H.: 1968, *Astrophys. J.* **154**, 891
 Peebles P.J.E.: 1980, in *The Large-Scale Structure of the Universe*, Princeton Univ. Press.
 Pfenniger D., Combes F.: 1994, *Astron. Astrophys.* **285**, 94
 Pfenniger D., Combes F., Martinet L.: 1994, *Astron. Astrophys.* **285**, 79
 Press W.H., Schechter P.: 1974, *Astrophys. J.* **187**, 425
 Puy D., Signore M.: 1996, *Astron. Astrophys.* **305**, 371
 Reimers D., Kohler S., Wisotzki L. et al.: 1997, *Astron. Astrophys.* **327**, 890
 Scalo J.M.: 1985, in *Protostars and Planets II*, ed. D.C. Black & M.S. Matthews, Univ. of Arizona Press, Tucson, p. 201
 Shapiro P.R., Kang H.: 1987, *Astrophys. J.* **318**, 32
 Silk J.: 1968, *Astrophys. J.* **151**, 459
 Stancil P.C., Lepp S., Dalgarno A.: 1996, *Astrophys. J.* **458**, 401
 Tegmark M., Silk J., Blanchard A.: 1994, *Astrophys. J.* **420**, 484
 Tegmark M., Silk J., Rees M.J., Blanchard A., Abel T., Palla F.: 1997, *Astrophys. J.* **474**, 1
 Uehara H., Susa H., Nishi R., Yamada M., Nakamura T.: 1996, *Astrophys. J.* **473**, L95
 White S.D.M., Rees M.J.: 1978, *Mon. Not. R. Astr. Soc.* **183**, 341
 Yoneyama 1972, *Publ. Astron. Soc. Japan* **24**, 87